



# TOMATO

# SKYWAY 687



Suited for gas green or vine ripe production. Widely adaptable throughout the Southeast U.S. at the start of the crop remove ground shoots (those that form at or below cotyledons) and a maximum of 0-2 axillary shoots.

Relative Days (TP)	65-70
Type	Hybrid Round Red Tomato
Fruit Size	Extra-Large to Jumbo
Fruit Shape	Globe
Plant Characteristics	Determinate
Resistance/Tolerance	HR: Fol: 0-2, ToMarV, Va: 0, Vd: 0 IR: Ma, Mi, Mj, TSWV, TYLCV

#### Pathogen and Pest Codes:

Fol	Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> )
Ma	Root-knot ( <i>meloidogyne arenaria</i> )
Mi	Root-knot ( <i>meloidogyne incognita</i> )
Mj	Root-knot ( <i>meloidogyne javanica</i> )
ToMarV	Tomato Marchitez Virus
TSWV	Tomato Spotted Wilt Virus
TYLCV	Tomato Yellow Leaf Curl Virus
Va	Verticillium Wilt ( <i>verticillium albo-atrum</i> )
Vd	Verticillium Wilt ( <i>verticillium dahliae</i> )

Stokes Seeds Ltd., 296 Collier Rd S., PO Box 10  
Thorold, ON L2V 5E9 Canada  
[www.StokeSeeds.com](http://www.StokeSeeds.com) / 800.263.7233

Stokes Seeds Inc., 2495 Walden Ave., PO Box 548  
Buffalo, NY 14042 USA  
[www.StokeSeeds.com](http://www.StokeSeeds.com) / 800.962.4999

Note: All variety information presented herein is based on field and laboratory observation. Actual crop yield and quality are dependent upon many factors beyond our control and NO WARRANTY is made for crop yield and quality. Since environmental conditions and local practices may affect variety characteristics and performance, we disclaim any legal responsibility for these. Read all tags and labels.They contain important conditions of sale, including limitations of warranties and remedies.